AD	)	

Award Number: DAMD17-03-1-0570

TITLE: Development of a Mouse Model for Determination of the Role of the Catechol Metabolites of Estradiol in Mammary Tumorigenesis

PRINCIPAL INVESTIGATOR: James D. Yager, Ph.D.

CONTRACTING ORGANIZATION: Johns Hopkins University

Baltimore, MD 21205-2179

REPORT DATE: August 2005

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20060525024

# REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arilington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently

4302. Respondents should b	e aware that notwithstanding an	y other provision of law, no perso	n shall be subject to any penalty	for failing to comply with	h a collection of information if it does not display a currently			
1. REPORT DATE		2. REPORT TYPE		3. [	DATES COVERED			
01-08-2005		<u>Final</u>			1 Jul 2003 –20 Jul 2005			
4. TITLE AND SUBTI					CONTRACT NUMBER			
Development of a Mouse Model for Determination of the Role of the Catechol								
Metabolites of Est	radiol in Mammary	Tumorigenesis			GRANT NUMBER			
				L	MD17-03-1-0570			
					PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)				5d.	PROJECT NUMBER			
James D. Yager, I	Ph.D.			5e.	TASK NUMBER			
				5f. WORK UNIT NUMBER				
	GANIZATION NAME(S)	AND ADDRESS(ES)	11.00		PERFORMING ORGANIZATION REPORT NUMBER			
Johns Hopkins University Baltimore, MD 21205-2179								
	I Research and Ma	NAME(S) AND ADDRES teriel Command	S(ES)	10.	SPONSOR/MONITOR'S ACRONYM(S)			
T OIL DOLLON, Mary	Idiid 21702 0012			1	SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited								
13. SUPPLEMENTAR	Y NOTES							
14. ABSTRACT								
model to study the genetic crossing, i the absence of CC resulting COMTKC were on-hand. Un began. Because of	e role of estrogen cantroduce the COMDMT on several biodo/ERCO/Wnt-1 femanticipated difficultiff the delay, a 1 year	atechol metabolites FKO genotype into to chemical end points ale mice. ERKO/Wr es were encountere or extension of the p	in mammary tumorion to the ERCO/Wnt-1 mident and on mammary on the control of the	genesis. The a be; 2) Initiate s gland developmed after notifing pairs were 2005 without a	ceptor/Wnt-1 (ERKO/Wnt-1) mouse aims were: 1) Through a complex studies to determine the effects of ment and tumorigenesis in the ication of award; COMTKO mice obtained and successful breeding additional funds was requested and generate the mice needed for the			
15. SUBJECT TERMS								
Breast Cancer *								
16. SECURITY CLASS		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC				
a. REPORT	b. ABSTRACT	c. THIS PAGE	1		19b. TELEPHONE NUMBER (include area			
U	U	υ	טט	6	code)			

# **Table of Contents**

Cover1
SF 2982
Table of Contents3
Introduction4
Body4
Key Research Accomplishments5
Reportable Outcomes5
Conclusions5
References5
Appendicesnonc

#### Introduction

The main risk factors associated with the development of breast cancer implicate estradiol (E2) as a causative agent<sup>1</sup>. The accepted hypothesis is that prolonged exposure to estrogen leads to persistent enhanced proliferation mediated through the binding of E2 to the estrogen receptor. However, E2 also undergoes cytochrome P450 (CYP)-mediated oxidative biotransformation to catechol metabolites, 2hydroxy (2-OH) and 4-hydroxy (4-OH) E2<sup>2</sup>. 4-OH E2 in particular has been shown to cause/contribute to estrogen carcinogenesis through further oxidative metabolism to quinones that form adducts with adenine and guanine in DNA and cause oxidative DNA damage through participation in redox cycling processes<sup>2</sup>. These catechol estrogen metabolites are present in mouse mammary gland tissue at concentrations of pmoles/mg tissue<sup>3</sup>. CEs are primarily inactivated by O-methylation catalyzed by catechol-O-methyltransferase (COMT)<sup>4</sup>. Inhibition of COMT enhanced E2-induced renal tumorigenesis in the Syrian hamster model<sup>4</sup> and DNA damage in human MCF-7 cells<sup>5</sup>, suggesting that COMT is highly protective against adverse effects caused by the catechol metabolites of E2. COMT is polymorphic in humans and several, but not all, epidemiology studies have observed that the genotype encoding a low activity COMT is associated with an increased risk for developing breast cancer in certain women<sup>6-8</sup>. These observations implicate the catechol metabolites of E2 in breast cancer causation. Experimental investigation of the extent and mechanisms of contribution of the estrogen catechol metabolites would be greatly facilitated by the availability of an appropriate animal model. Mammary tumor incidence in intact estrogen receptor knockout (ERKO)/Wnt-1 female mice is 20%, whereas it is reduced to 8% in ovariectomized females 9. These results demonstrated that in the absence of ER expression, estrogen still contributes to mammary tumor formation. Our hypothesis regarding the role of estrogen in these mice is that the catechol metabolites undergo oxidative biotransformation to quinones which then form DNA adducts directly, or oxidative DNA damage indirectly through redox cycling, and subsequently mutations, which contribute to mammary tumor formation. Since COMT is protective, the absence of COMT would be expected to increase the incidence and/or shorten the latency of tumorigenesis in the ERKO/Wnt-1 mice. Thus, the purpose of the research proposed for this Concept Grant was to develop a catechol-o-methyltransferase knockout (COMTKO)-estrogen receptor/Wnt-1 (ERKO/Wnt-1) mouse model for use in studies on the role of estrogen catechol metabolites in mammary tumorigenesis. The aims were: 1) Through a complex genetic crossing, introduce the COMTKO genotype into the ERCO/Wnt-1 mice; 2) Initiate studies to determine the effects of the absence of COMT on several biochemical end points reflecting the effects of the absence of COMT (e.g. estrogen catechol metabolite levels, DNA damage) and on mammary gland development and tumorigenesis in COMTKO/ERCO/Wnt-1 female mice.

## **Body**

Preliminary studies, partly supported by this Concept grant, revealed that mammary tissue from COMTKO mice formed catechol metabolites of E2, but as expected, not the methoxy metabolites whch are formed when COMT is expressed. Furthermore, oxidative DNA damage in COMTKO mice was increased compared to the level observed in COMT wild type mice (See Figure 1).

After notification of award, ERKO/Wnt-1 mice, in a C57Bl/6 background, were obtained from Dr. Richard Santen (University of Virginia); the C57Bl/6 COMTKO mice were on-hand. However, unanticipated breeding difficulties were encountered. The COMTKO mice went through a period where they stopped breeding. Additional breeding pairs were obtained from the originator of the strain at Rockefeller University. By the time the mice were cleared through quarantine and breeding could commence, the project was about 6 months behind schedule. Thus, a request was made and granted for an extension of the project for a second year to August 20, 2005without additional funds.

The breeding scheme is complex. The genotype of the female mice needed to test the hypothesis is catechol-O-methyltransferase knockout/estrogen receptor knockout/Wnt (COMTKO/ERKO /Wnt).

However, ERKO females and males are infertile and Wnt females cannot nurse their young. Thus, the desired females must be obtained from the following breeding pairs: Female COMTKO/ER<sup>+/-</sup>; Male – COMTKO/ER<sup>+/-</sup>/Wnt (Wnt is carried in the males as a single copy).

## **Key Research Accomplishments**

- 1. The COMTKO genotype was introduced into female ER<sup>+/-</sup> and male ER<sup>+/-</sup>/Wnt<sup>+/-</sup> mice to generate breeding pairs needed to produce mice of the desired genotype. Seven breeding pairs, which are **Female:** COMTKO/ER<sup>+/-</sup>; **Male:** COMTKO/ER<sup>+/-</sup>/Wn<sup>+/-</sup>, are now in place.
- 2. Breeding is ongoing. It was expected that  $1/16^{th}$  of the pups would be females with the desired genotype of COMTKO/ERKO/Wnt. However, to date, of the 57 pups genotyped, while approximately 3 were expected to be COMTKO/ERKO/Wnt, none have been obtained. Breeding is continuing. Control female mice that are ERKO/Wnt are also being bred.

## Reportable Outcomes

None at this time, other than the fact that the desired breeding pairs have been developed.

#### Conclusion

No conclusions can be drawn at this point. Unexpected problems were encountered in breeding the COMTKO mice that caused an initial delay in progress. Once new breeding pairs of COMTKO mice were obtained, breeding began to develop the breeding pairs needed to produce female mice of the genotype needed. In this sense, the Concept Project has been successful, since the primary goal of developing the desired breeding pairs has been accomplished. It will now be a matter of time for sufficient number (minimum, 30) of COMTKO/ERKO/Wnt female mice to be identified and maintained for 15-18 months in order to determine the effect of the absence of COMT on mammary tumor incidence.

#### References

- 1. Key T, Appleby P, Barnes I, Reeves G. Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. J Natl Cancer Inst 2002;94(8):606-16.
- 2. Cavalieri EL, Rogan EG. A unifying mechanism in the initiation of cancer and other diseases by catechol quinones. Ann N Y Acad Sci 2004;1028:247-57.
- Devanesan P, Santen RJ, Bocchinfuso WP, Korach KS, Rogan EG, Cavalieri E. Catechol estrogen metabolites and conjugates in mammary tumors and hyperplastic tissue from estrogen receptor-alpha knock-out (ERKO)/Wnt-1 mice: implications for initiation of mammary tumors. Carcinogenesis 2001;22(9):1573-6.
- 4. Monographs JNCI. Estrogens as endogenous carcinogens in the breast and prostate. JNCI Monographs 2000;27:159.

- 5. Lavigne JA, Goodman JE, Fonong T, et al. The effects of catechol-O-methyltransferase inhibition on estrogen metabolite and oxidative DNA damage levels in estradiol-treated MCF-7 cells. Cancer Res 2001;61(20):7488-94.
- 6. Lavigne JA, Helzlsouer KJ, Huang HY, et al. An association between the allele coding for a low activity variant of catechol-O-methyltransferase and the risk for breast cancer. Cancer Res 1997;57(24):5493-7.
- 7. Miyoshi Y, Noguchi S. Polymorphisms of estrogen synthesizing and metabolizing genes and breast cancer risk in Japanese women. Biomed Pharmacother 2003;57(10):471-81.
- 8. Mitrunen K, Hirvonen A. Molecular epidemiology of sporadic breast cancer. The role of polymorphic genes involved in oestrogen biosynthesis and metabolism. Mutat Res 2003;544(1):9-41.
- 9. Yue W, Santen RJ, Wang JP, et al. Genotoxic metabolites of estradiol in breast: potential mechanism of estradiol induced carcinogenesis. J Steroid Biochem Mol Biol 2003;86(3-5):477-86.

**Figure** 

# Oxidative DNA Damage in COMTKO vs WT Mice

